Table 2 b: Extensive list of well-known natural compounds (drugs, lead compounds for drugs or compounds in clinical trials)) used in areas of neurological diseases, immunological or inflammatory processes, cardiovascular diseases and oncological diseases and similar compounds (Tanimoto >= 0.85) from the SuperNatural Database

* Anatomical Therapeutic Chemical (ATC) classification code generated by the World Health Organization (WHO) describes the therapeutic subgroup (1)

| Natural Compound | Status (Reference) | Similar Compounds in SuperNatural (Tanimoto >=85%) | Analytical Therapeutic Chemical (ATC) classification code* |
|----------------------|--|--|--|
| | Neurological disease area | | |
| Physostigmine | Lead compound of Eserine Sulfate (2) | 1 | S01EB05* (Parasympathomimetics) V03AB* (Antidotes) |
| Huperzine | Phase III clinical trials (3) | 1 | V03* (All other therapeutic products) |
| Morphine | Lead compound of Roxanol TM (2) | 9 | N02AA* (Natural opium alkaloids) |
| Tetrahydrocannabinol | Lead compound of Tetrahydrocannabinol (2) | 2 | A04AD* (Other antiemetics) |
| | Cardiovascular | | |
| Erythromycin | Phase II clinical trials (2) | 25 | J01FA* (Macrolides) S01AA* (Antibiotics) |
| | Immunological,inflammatory | | |
| Cyclosporine | Lead compound of Sandimmune® (2) | 3 | L04AA* (Selective immunosuppressive agents) |
| Tacrolimus | Lead compound of Protopic® (2) | 2 | D11AX* (Other dermatologicals) L04AA* (Selective immunosuppressive agents) |
| Sirolimus | Lead compound of Rapamune® (2) | 2 | L04AA* (Selective immunosuppressive agents) |
| | Oncological disease area | | |
| Camptothecin | Phase I & II & III clinical trials (2) | 7 | L01* (Antineoplastic agents) |

| Homoharringtonine | Phase II clinical trials | 1 | L01* (Antineoplastic agents) |
|-------------------|------------------------------------|----|--|
| | (2) | | |
| Paclitaxel | Lead compound of Taxol® | 5 | L01CD* (Taxanes) |
| | (4) | | |
| Protopanaxadiol | Phase I clinical trials | 17 | L01* (Antineoplastic agents) |
| | (2) | | |
| Triptolide | Phase I clinical trials | 18 | L01* (Antineoplastic agents) |
| _ | (2) | | |
| Vinblastine | Lead compound of Velban® | 9 | L01CA* (Vinca alkaloids and analogues) |
| | (4) | | |
| Doxorubicin | Lead compound of Adriblastin® | 12 | L01DB* (Anthracyclines and related |
| | (5) | | substances) |
| Staurosporine | Phase I & II & III clinical trials | 4 | L01* (Antineoplastic agents) |
| - | (2) | | |
| Fumagillin | Lead compound of CKD-732 | 1 | L01* (Antineoplastic agents) |
| | (6) | | |
| Halichondrin B | Phase I clinical trials | 5 | L01* (Antineoplastic agents) |
| | (2) | | |

- 1. WHO. (2003) World Health Organ Tech Rep Ser, **914**, 1-126.
- 2. Butler, M.S. (2005) *Nat Prod Red*, **22**, 162-195.
- 3. (2004) *Drugs R D*, **5**, 44-45.
- 4. Gabai, V., Budagova, K. and Sherman, M. (2005) Oncogene, 24, 3328-3338.
- 5. Koivusalo, R., Krausz, E., Helenius, H. and Hietanen, S. (2005) *Mol Pharmacol.*, **68**, 372-382.
- 6. Kim, E. and Shin, W. (2005) *Biol Pharm Bull*, **28**, 217-223.